

FARMACI CON USO CONSOLIDATO NEL TRATTAMENTO CORRELATO AI TRAPIANTI PER INDICAZIONI ANCHE DIFFERENTI DA QUELLE PREVISTE DAL PROVVEDIMENTO DI AUTORIZZAZIONE ALL'IMMISSIONE IN COMMERCIO

BASILIXIMAB

Indicazioni già autorizzate:

• Indicato per la profilassi del rigetto acuto in pazienti adulti e pediatrici sottoposti a trapianto renale allogenico de novo. Deve essere usato in associazione ad un trattamento immunosoppressivo a base di ciclosporina in microemulsione e corticosteroidi nei pazienti con una quantità di anticorpi reattivi inferiore all'80%, o in uno schema terapeutico immunosoppressivo di mantenimento in triplice terapia comprendente ciclosporina in microemulsione, corticosteroidi e azatioprina o micofenolato mofetile.

Estensione delle indicazioni:

- Profilassi del rigetto acuto in pazienti adulti e pediatrici sottoposti a trapianto di fegato e a Trapianto isole di Langerhans.
- Profilassi del rigetto acuto in pazienti adulti sottoposti a trapianto di rene e pancreas¹⁻²⁰.
- Profilassi del rigetto acuto in pazienti adulti e pediatrici sottoposti a trapianto di cuore²¹.

EVEROLIMUS

Indicazioni già autorizzate:

• Profilassi del rigetto d'organo in pazienti adulti, a rischio immunologico da lieve a moderato, sottoposti a trapianto renale o cardiaco allogenico. Certican deve essere utilizzato in associazione con ciclosporina in microemulsione e corticosteroidi.

Estensione delle indicazioni:

- Profilassi del rigetto acuto in pazienti pediatrici sottoposti a trapianto di rene.
- Profilassi del rigetto acuto in pazienti adulti sottoposti a trapianto di polmone.
- Profilassi del rigetto acuto in pazienti adulti sottoposti a trapianto di fegato.
- Pazienti con trapianto di fegato che necessitano la riduzione/sospensione nell'inibitore della calcineurina con problemi di tossicità renale²²⁻³⁰.
- Trapianto di fegato pediatrico³¹.

Fattori di crescita dei leucociti (filgrastim, lenograstim)

Indicazioni già autorizzate:

• Trattamento della neutropenia

Estensione delle indicazioni:

• Neutropenia (neutrofili < 750/L) nei pazienti trapiantati di fegato o con diagnosi clinica di cirrosi, che ricevono interfe-

rone standard o peghilato in monoterapia o in combinazione con ribavirina e che presentano risposta virologica precoce alla terapia³²⁻³⁹.

Immunglobulina di coniglio antitimocitaria

Indicazioni già autorizzate:

• Profilassi nell'adulto della malattia acuta e cronica da trapianto verso ospite in trapianti da donatore familiare non immunocompatibile o immunocompatibile non familiare (*Graft versus Host Disease, GvHD*).

Estensione delle indicazioni:

- Trattamento nell'adulto della malattia acuta e cronica da trapianto verso ospite (*Graft versus Host Disease, GvHD*).
- Profilassi e trattamento nel paziente pediatrico della malattia acuta e cronica da trapianto verso ospite (*Graft versus Host Disease, GvHD*).
- Regime di condizionamento nel trapianto autologo per malattie autoimmuni⁴⁰⁻⁴⁷.

Micofenolato Mofetile

Indicazioni già autorizzate:

- Trapianto rene.
- Trapianto rene pediatrico.
- Trapianto cuore.
- Trapianto fegato.

Estensione delle indicazioni:

- Trapianto cuore pediatrico;
- Trapianto fegato pediatrico;
- Trapianto pancreas
- Trapianto polmone
- Trapianto di midollo osseo
- Trapianto isole di Langerhans⁴⁸⁻⁵².

Micofenolato Sodico

Indicazioni già autorizzate:

• Myfortic è indicato in associazione con ciclosporina e corticosteroidi, per la profilassi del rigetto acuto, in pazienti adulti che ricevono un trapianto allogenico di rene.

Estensione delle indicazioni:

- Profilassi del rigetto acuto in pazienti adulti con trapianto di cuore. Pazienti adulti con trapianto di fegato e di cuore in cui il micofenolato mofetile dia effetti collaterali di tipo gastroenterico che richiedano la diminuzione/sospensione della dose⁵³⁻⁵⁹.
- Profilassi del rigetto acuto nel trapianto di rene in associazione con Tacrolimus⁶⁰.

RAPAMICINA

Indicazioni già autorizzate:

- Trapianto rene.

Estensione delle indicazioni:

- Trapianto fegato;
- Trapianto pediatrico di fegato e/o rene.
- Trapianto midollo.
- Trapianto pancreas, cuore, polmone
- Trapianto isole di Langerhans⁶¹⁻⁶⁵.

TACROLIMUS

Indicazioni già autorizzate:

- Profilassi AR e trattamento AR in: trapianto rene, trapianto fegato, trapianto cuore; trattamento AR in: trapianto rene-pancreas, trapianto pancreas, trapianto cuore-polmone, trapianto polmone, trapianto intestino multi viscerale.

Estensione delle indicazioni:

- Profilassi AR in trapianto di cuore-polmone, trapianto polmone, trapianto pancreas, trapianto rene-pancreas, trapianto intestino trapianto isole di Langerhans.
- Profilassi AR e trattamento GVHD in trapianto midollo osseo⁶⁶⁻⁷⁰.
- Tutte le indicazioni anche in associazione con altri farmaci ad attività immunosoppressiva o immunomodulante e/o steroide⁷¹.

THYMOGLOBULINE (ATC: L04AA04)

Indicazioni già autorizzate:

- Profilassi nell'adulto della malattia acuta e cronica da trapianto verso ospite (Graft vs Host Disease) nel trapianto da donatore familiare non immunocompatibile o immunocompatibile non familiare.

Estensione delle indicazioni:

- Profilassi nell'adulto della malattia acuta e cronica da trapianto verso ospite (GvHD) nel trapianto da donatore familiare immunocompatibile (HLA identico)⁷²⁻⁷⁹.

VALGANCICLOVIR

Indicazioni già autorizzate:

- Prevenzione della malattia da CMV in pazienti CMV negativi e sottoposti a trapianto d'organo SOLIDO da donatore CMV positivo. trattamento di induzione e mantenimento della retinite da CMV in pazienti con AIDS.

Estensione delle indicazioni:

- Trapianto midollo osseo;
- Trapianto d'organo solido pediatrico^{80,81}.

BIBLIOGRAFIA

1. Spada M et al. Randomized Trial of Basiliximab Induction versus steroid Therapy in Pediatric Liver Allograft Recipients Under Tacrolimus Immunosuppression Am J Transplantation 2006; 6: 1913-1921. **2.** Gruttadauria S et al. A Safe Immunosuppressive Protocol in Adult to-Adult Living Related Liver Transplantation. Transplant Proc 38, 1106-1108 (2006). **3.** Lladó L et al. Immunosuppression without steroids in liver transplantation is safe and reduces infection and metabolic

complications: Results from a prospective multicenter randomized study. J. Hepatol 44 (2006) 710-718. **4.** Orr DW et al. Anti-Interleukin 2 Receptor Antibodies and Mycophenolate Mofetil for Treatment of Steroid - Resistant Rejection in Adult Liver Transplantation. Transplant Proc 37, 4373-4379 (2005). **5.** Ganschow R et al. Long-term results of basiliximab induction immunosuppression in paediatric liver transplants recipients. Pediatr Transplantation 2005; 9: 741-745. **6.** Spada M et al. An anti-interleukin 2 receptor monoclonal antibody to reduce the incidence of acute cellular rejection after liver transplantation. Pediatr Transplantation 2000, Vol 4., p. 62 (Abs P49). **7.** Ganschow R et al. The anti-interleukin 2 receptor antibody basiliximab after pediatric liver transplantation: a pilot study. Pediatr Transplantation 2000, Vol. 4, p. 95 (Abs O145). **8.** Ganschow R et al. First experience with basiliximab in pediatric liver graft recipients. Pediatr Transplantation 2001; 5: 353-358. **9.** Filippini F et al. Study of Simulect-Based, Steroid-Free Immunosuppressive Regimen in HCV+ De Novo Liver Transplant Patients: Preliminary Results. Transplantation Proceedings 2001; 33: 3211-12. **10.** Kelly DA. The use of anti-interleukin 2 receptor antibodies in pediatric liver transplantation. Pediatr Transplantation 2001; 5: 386-389. **11.** Neuhaus P et al. Improved Treatment Response With Basiliximab Immunoprophylaxis After Liver Transplantation: Results from a Double-Blind Randomized Placebo-Controlled Trial. Liver Transplantation 2002; 8, 132-142. **12.** Cantarovich M et al. Anti-CD25 Monoclonal antibody coverage allows for calcineurin inhibitor "Holiday" in solid organ transplant patients with acute renal dysfunction. Transplantation Vol 73 n 7. **13.** Nashan B. The Interleukin "Pathway" and the Route to Logical Immunosuppression. Current Issues in Liver and Small Bowel Transplantation 2002, 9, 164-174. **14.** Kuse E et al. Immunoprophylaxis with Simulect® (Basiliximab) in combination with Cyclosporine and Steroids in Liver Transplantation. American Journal of Transplantation 2001, vol 1, p 202 (Abs 266). **15.** Venze A et al. Basiliximab monotherapy following B-cell lymphoma after pediatric liver transplantation and anti- CD20 therapy. Pediatr Transplantation 2003; 7: 404-407. **16.** Reding R et al. Steroid-free liver transplantation in children. Lancet 2003; 362: 2068-70. **17.** Liu CL et al. Interleukin 2-Receptor Antibody (Basiliximab) for Immunosuppressive Induction Therapy After Liver Transplantation: A Protocol With Early Elimination of Steroids and Reduction of Tacrolimus Dosage. Liver Transplantation, 10: No. 6 (June) 728-733. **18.** Gruttadauria S et al. Basiliximab in a Triple-Drug Regimen with Tacrolimus and Steroids in Liver Transplantation. Transplantation Proceedings, 37, 2611-2613 (2005). **19. Trapianto di isole di Langerhans.** Oberholzer J, Toso C, Triponez F, Ris F, Bucher P, et al: Human islet allotransplantation with Basiliximab in type I diabetic patients with end-stage renal failure. Transplantation Proceedings, 34, (3) 823-825. **20. Trapianto di Rene-Pancreas.** Zhang R, Florman S, Devidoss S, Zarifian A, Yau CL, Paramesh A, Killackey M, Alper B, Fonseca V, Slakey D. A comparison of long-term survivals of simultaneous pancreas-kidney transplant be-

tween African American and Caucasian recipients with basiliximab induction therapy. *Am J Transplant*. 2007 Jul;7(7):1815-21.

Boggi U, Vistoli F, Del Chiaro M, Signori S, Amoreso G, Vahadia Bartolo T, Sgambelluri F, Barsotti M, Tregnaghi C, Paleologo G, Coppelli A, Giannarelli R, Rizzo G, Marchetti P, Mosca F.

Neoral versus prograf in simultaneous pancreas-kidney transplantation with portal venous drainage: three-year results of a single-center, open-label, prospective, randomized pilot study. *Transplant Proc*. 2005 Jul-Aug;37(6):2641-3.

Chow FY, Polkinghorne K, Saunder A, Kerr PG, Atkins RC, Chadban SJ. Historical controlled trial of OKT3 versus basiliximab induction therapy in simultaneous pancreas-renal transplantation. *Nephrology (Carlton)*. 2003 Aug;8(4):212-6.

21. Pape L et al. Reversal of loss of glomerular filtration rate in children with transplant nephropathy after switch to everolimus and low-dose cyclosporine A. *Pediatr Transplantation* 2007; 11: 291-293.

22. Vester U et al. Everolimus (Certican) in combination with Neoral in Pediatric Renal Transplant Recipients: Interim analysis after 3 Months. *Transplantation Proceedings* 2002; 34, 2209-22.

23. Hoyer P et al. Everolimus in Pediatric de Novo Renal Transplant Patients. *Transplantation* 2003; 75: 2082-2085.

24. Van Damme-Lombaerts R et al. Single-dose pharmacokinetics and tolerability of everolimus in stable pediatric renal transplant patients. *Pediatric Transplantation* 2002; 6: 147-152.

25. Kovarik JM et al. Everolimus in Pulmonary Transplantation: Pharmacokinetics and Exposure-Response Relationship. *The Journal of Heart and Lung Transplantation* 2006; 25: No 4.

26. Everolimus versus Azathioprine in Maintenance Lung Transplant Recipients: An International, Randomized Double-Blind Clinical Trial. *American Journal of Transplantation* 2006; 6: 169-177.

27. Azzola et al. Everolimus and Mycophenolate Mofetil are potent inhibitors of fibroblast proliferation after lung Transplantation. *Transplantation Vol 77, No. 4* (2004).

28. Everson G. Everolimus and mTOR Inhibitors in Liver Transplantation: Opening the "Box". *Liver Transplantation* 12: 1571-1573, 2006.

29. Poirier CD. Promise of Neoral C2; Basiliximab and Everolimus in Lung Transplantation. *Transplantation Proceedings*, 36 (Suppl 2S)N, 5095-5135 (2004).

30. Levy e al. Safety, Tolerability and Efficacy of Everolimus in De Novo Liver Transplant Recipients: 12-and 36-Month Results. *Liver Transplantation* 12: 1640-1648, 2006.

31. **Trapianto di fegato.** Katrina A. Ford *Paediatric Immunology Following Solid Organ Transplantation Arch Dis Child Educ Ed* 2006; 91:87-91.

32. Manns MP et al. and the International Hepatitis Interventional Therapy Group. Peginterferon alfa-2b plus ribavirin compared with interferon alfa-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomized trial. *Lancet* 2001;358:958-65.

33. Fried MW et al. Peg Interferon alf 2a plus Ribavirin in chronic hepatitis C virus infection. *N Engl J Med* 2002; 347: 975-82.

34. Higashi Y et al. Case report: agranulocytosis induced by interferon alpha therapy for chronic hepatitis C *J Gastroenterol Hepatol* 1996; 11:1012-1015.

35. Van Thiel DH et al. Combination treatment of advanced HCV associated liver disease with interferon and

GCSF. *Hepatogastroenterology* 1995; 42:907-12.

36. Fukuda A et al. Effects of interferon alpha on peripheral neutrophil counts and serum granulocyte colony-stimulating factor for the treatment of chronic hepatitis C *Cell Mol Ther* 2000; 6:149-154.

37. Carreno V et al. Randomized controlled trial of recombinant human granulocyte-macrophage colony-stimulating factor for the treatment of chronic hepatitis C *Cytokine* 2000; 12: 165-70.

38. Shiffman ML et al. Use of granulocyte macrophage colony stimulating factor alone or in combination with interferon-alpha-2b for treatment of chronic hepatitis C *J Hepatol* 1998; 28: 382-89.

39. National Institutes of health consensus development conference statement: Management of hepatitis C: 2002 –June 10-12 2002. *Hepatology* 2002; 36: S3-S20.

40. Hsu B et al. Use of antithymocyte globulin for treatment of steroid-refractory acute graft-versus-host disease: an international practice survey. *Bone Marrow Transplantation* 2001; 28: 945–950.

41. Graziani F et al. Treatment of acute graft-versus-host disease with low-dose, alternate-day antithymocyte globulin *Haematol* 2002; 87: 973-978.

42. Bacigalupo A et al. Fludarabine, cyclophosphamide and anti-thymocyte globulin for alternative donor transplants in acquired severe aplastic anemia: a report from the EBMT-SAA Working Party. For the Severe Aplastic Anemia Working Party of the European Group for Blood and Marrow Transplantation (SAA WP-EBMT) *Bone Marrow Transplantation* 2005; 36: 947–950.

43. Kang HJ et al. Fludarabine, cyclophosphamide plus thymoglobulin conditioning regimen for unrelated bone marrow transplantation in severe aplastic anemia. *Bone Marrow Transplantation* 2004; 34: 939–943.

44. Horn B et al. Reduced intensity conditioning using intravenous busulfan, fludarabine and rabbit ATG for children with nonmalignant disorders. *Bone Marrow Transplantation* 2006; 37: 263–269.

45. Kikuta A. Nonmyeloablative stem cell transplantation for nonmalignant diseases in children with severe organ dysfunction. *Bone Marrow Transplantation* 2006; 38: 665–669.

46. Hsu B et al. Use of antithymocyte globulin for treatment of steroid-refractory acute graft-versus-host disease: an international practice survey. *Bone Marrow Transplantation* 2001; 28: 945–950.

47. Tagliabue A et al. Favourable response to antithymocyte globulin therapy in resistant acute graft-versus-host disease. *Bone Marrow Transplantation* 2005; 36: 45.

48. **Trapianto di cuore pediatrico.** J. Agüero et al. Influence of Immunosuppression Regimen on Heart Transplantation Survival. *Transplantation Proceedings* 2006; 38: 2550–2; Olivia Boyer et al. Improvement of Renal Function in Pediatric Heart Transplant Recipients Treated with Low-Dose Calcineurin Inhibitor and Mycophenolate Mofetil. *Transplantation* 2005; 79 (10): 1405 1410; Dipchand AI et al. Mycophenolate mofetil in pediatric heart transplant recipients: A single-center experience. *Pediatr Transplantation* 2001; 5: 112–118; Tonshoff B et al. Treatment strategies in pediatric solid organ transplant recipients with calcineurin inhibitor-induced nephrotoxicity. *Pediatr Transplantation* 2006; 10: 721–729; Groetzner J et al. Cardiac transplantation in pediatric pa-

tients: fifteen-year experience of a single center. *Ann Thorac Surg* 2005; 79 (1): 53-60; *Since the introduction of mycophenolate mofetil, freedom from acute rejection increased to 62%;* Gajarski RJ et al. Lack of correlation between MMF dose and MPA level in pediatric and young adult cardiac transplant patients: does the MPA level matter? *Am J Transplant* 2004; 4 (9): 1495-500; Shaddy RE et al. Mycophenolic mofetil reduces the HLA antibody response of children to valved allograft implantation. *Ann-Thorac-Surg* 2004; 77 (5): 1734-9. *This study demonstrates the ability to pharmacologically abrogate the HLA class I antibody response to valved allograft implantation in children using MMF.* Boucek RJ, Boucek MM Pediatric heart transplantation. *Curr-Opin-Pediatr* 2002; 14 (5): 611 9. *Currently, recipients are maintained on immunosuppressive medications that target calcineurin (eg, cyclosporine, tacrolimus), lymphocyte proliferation (eg, azathioprine, mycophenolate mofetil (MMF), sirolimus) and, in some instances antiinflammatory corticosteroids.* Kobashigawa Review of Major Clinical Trials with Mycophenolate Mofetil in Cardiac Transplantation. *Transplantation* 2005; 80 (25): S235-S243. **49. Trapianto di Fegato pediatrico.** Renz JF et al. Mycophenolate mofetil, microemulsion cyclosporine, and prednisone as primary immunosuppression for pediatric liver transplant recipients. *Liver Transpl Surg.* 19995(2): 136-43. Chardot C et al. Use of mycophenolate mofetil as rescue therapy after pediatric liver transplantation. *Transplantation* 2001; 71 (2): 224-9 *These preliminary results suggest that MMF is an effective and safe immunosuppressant in pediatric LT recipients. Its use is hampered by frequent gastrointestinal and hematological side-effects. MMF does not seem to increase the risk of PTLN nor CMV disease.* Aw MM et al. Calcineurin-inhibitor related nephrotoxicity- reversibility in paediatric liver transplant recipients. *Transplantation* 2001; 72 (4): 746-9. *MMF allows the recovery of renal function from CI related nephrotoxicity in more than 70% of paediatric liver transplant recipients with renal impairment.* Ferraris JR et al. Mycophenolate mofetil and reduced doses of cyclosporine in pediatric liver transplantation with chronic renal dysfunction: changes in the immune responses. *Pediatr Transplant* 2004; 8 (5): 454. Evans HM et al. Mycophenolate Mofetil for Renal Dysfunction after Pediatric Liver Transplantation. *Transplantation* 2005; 79: 1575-1580. Klupp J et al. Indications of Mycophenolate Mofetil in Liver Transplantation. *Transplantation* 2005; 80: S142-S14. Marion M. Mycophenolic Acid Pharmacokinetics in Pediatric Liver Transplant Recipients. *Liver Transplantation* 2003; 9 (4): 383-388. Apezato ML et al. Mycophenolate mofetil promotes prolonged improvement of renal dysfunction after pediatric liver transplantation: Experience of a single center. *Pediatr Transplantation* 2007; 11: 82-86. Tredger JM. Monitoring Mycophenolate in Liver Transplant Recipients: Toward a Therapeutic Range. *Liver Transplantation* 2004; 10 (4): 492-502. Mention K et al. Renal function outcome in pediatric liver transplant recipients. *Pediatr-Transplant* 2005; 9 (2): 201-7. Nobili V et al.

Mycophenolate mofetil in pediatric liver transplant patients with renal dysfunction: preliminary data. *Pediatr-Transplant* 2003; 7 (6): 454-7. **50. Trapianto di Pancreas.** Odorico JS et al. Improved solitary pancreas transplant graft survival in the modern immunosuppressive era. *Cell-Transplant* 2000; 9 (6): 919-27. Rigotti P et al. Mycophenolate mofetil (MMF) versus azathioprine (AZA) in pancreas transplantation: a single-center experience. *Clin-Nephrol* 2000; 53 (4): 52-4. *In conclusion, patients treated with MMF required less frequent and less intensive treatment for acute rejection. However, its short- and long-term side effects should be further investigated.* Gruessner RW et al. Mycophenolate mofetil in pancreas transplantation. *Transplantation* 1998; 66 (3): 318-23 *For SPK recipients, the incidence of acute reversible rejection episodes was significantly lower with MMF than with azathioprine.* Paleologo G et al. Solitary pancreas transplantation: preliminary findings about early reduction of proteinuria in incipient or evident diabetic type I nephropathy. *Transplant Proc* 2004; 36 (3): 591-6. *The immunosuppressive therapy used basilixmab induction and tacrolimus, mycophenolate mofetil (MMF), and low dose steroid maintenance therapy.* Kaufman DB et al. Pancreas transplantation at Northwestern University. *Clin Transpl* 2000; 239-46. *We have found that multimodal immunotherapy including induction with tacrolimus-based maintenance combined with either MMF or sirolimus, with or without corticosteroids, resulted in excellent patient and graft survival rates with low rates of rejection.* Sutherland DE et al. Lessons learned from more than 1,000 pancreas transplants at a single institution. *Ann Surg* 2001; 233 (4): 463-501. *From December 16, 1966, to March 31, 2000, the authors performed 1,194 pancreas transplants. The analyses were divided into five eras: era 0, 1966 to 1973 (n = 14), historical; era 1, 1978 to 1986 (n = 148), transition to cyclosporine for immunosuppression, multiple duct management techniques, and only solitary (PAK and PTA) transplants; era 2, 1986 to 1994 (n = 461), all categories (SPK, PAK, and PTA), predominantly bladder drainage for graft duct management, and primarily triple therapy (cyclosporine, azathioprine, and prednisone) for maintenance immunosuppression; era 3, 1994 to 1998 (n = 286), tacrolimus and mycophenolate mofetil used; and era 4, 1998 to 2000 (n = 275), use of daclizumab for induction immunosuppression, primarily enteric drainage for SPK transplants, pretransplant immunosuppression in candidates awaiting PTA Patient and graft survival rates have significantly improved over time as surgical techniques and immunosuppressive protocols have evolved.* Jordan ML et al. Steroid withdrawal for pancreas transplants under tacrolimus immunosuppression. *Transplant- Proc* 2001; 33 (1-2): 1655 Egidì FM. Management of Hyperglycaemia After Pancreas Transplantation. Are New Immunosuppressants the Answer? *Drugs* 2005; 65 (2): 153-166 Garcia VD et al. Immunosuppression in Pancreas Transplantation: Mycophenolate Mofetil Versus Sirolimus. *Transplantation Proceedings* 2004; 36: 975-977 Gruessner AC, Sutherland DER. Pancreas transplant outcomes

for United States (US) and non-US cases as reported to the United Network for Organ Sharing (UNOS) and the International Pancreas Transplant Registry (IPTR) as of June 2004. Clin Transplant 2005; 19: 433-455. Jung M et al. Comparison of Azathioprine and Mycophenolate Mofetil for the Prevention of Acute Rejection in Recipients of Pancreas Transplantation. J Clin Pharmacol 2001;41: 861-869. *In conclusion, compared with AZA, MMF significantly reduces the rate of biopsy-proven pancreas rejection during the first 6 months of transplantation and is well tolerated.* Mark W et al. Impact of Steroid Withdrawal on Metabolic Parameters in a Series of 112 Enteric/Systemic-Drained Pancreatic Transplants. Transplantation Proceedings 2005, 37, 1821-1825. **51. Trapianto di Polmone** Izbicki G et al. Improved Survival After Lung Transplantation in Patients Treated With Tacrolimus/Mycophenolate Mofetil as Compared With Cyclosporine/Azathioprine. Transplantation Proceedings 2002; 34: 3258-3259. Lama R et al. Lung Transplants With Tacrolimus and Mycophenolate Mofetil: A Review. Transplantation Proceedings 2003; 35: 1968-1973 Palmer SM et al. Results of a randomized, prospective, multicenter trial of Mycophenolate Mofetil versus azathioprine in the prevention of acute lung allograft rejection. Transplantation 2001; 71: 1772-1776. McNeil K et al. Comparison of Mycophenolate Mofetil and Azathioprine for Prevention of Bronchiolitis Obliterans Syndrome in De Novo Lung Transplant Recipients. Transplantation 2006; 81 (7): 998-1003. Roman A et al. Preliminary Results of Rescue Therapy With Tacrolimus and Mycophenolate Mofetil in Lung Transplanted Patients With Bronchiolitis Obliterans. Transplantation Proceedings 2002; 34: 146-147. Treede H et al. Tacrolimus versus Cyclosporine after Lung Transplantation: A Prospective, Open, Randomized Two-Center Trial Comparing Two Different Immunosuppressive Protocols. J Heart Lung Transplant 2001; 20: 511-517. Zuckermann A et al. Comparison Between Mycophenolate Mofetil and Azathioprine-Based Immunosuppressions in Clinical Lung Transplantation. J Heart Lung Transplant 1999;18: 432-440. Zuckermann A et al. Benefit of Mycophenolate Mofetil in Patients With Cyclosporine A-Induced Nephropathy. Bhorade SM et al. Comparison of three tacrolimus-based immunosuppressive regimens in lung transplantation. Am J Transplant. 2003; 3 (12): 1570-5. Addition of daclizumab and MMF to a tacrolimus-based immunosuppressive regimen decreased the incidence of acute rejection episodes without increasing any adverse events in our lung transplantation population. Groetzner J et al. Conversion to sirolimus and mycophenolate can attenuate the progression of bronchiolitis obliterans syndrome and improves renal function after lung transplantation. Transplantation. 2006; 81 (3): 355-60. *After BOS was diagnosed, conversion to MMF and Sir stabilized graft function only in some of the converted patients. Therefore, earlier administration of Sir-based immunosuppression might be a more promising approach. Whether conversion to CNI-free immunosuppression can actually ameliorate the ex-*

tent or progression of BOS has to be investigated in randomized trials. **52. Trapianto di isole di Langerhans** Shapiro AMJ, Ricordi C, Hering BJ et al: International trial of the Edmonton Protocol for islet transplantation. N Engl J Med 2006; 355: 1318-30, Froud T, Baidal DA, Ponte G, Ferreira DV, Ricordi C, Alejandro R: Resolution of neurotoxicity and beta-cell toxicity in an islet transplant recipient following substitution of tacrolimus with MMF. Cell transplant 2006, 15 (7): 613-629. **53.** Kobashigawa J et al. Similar Efficacy and Safety of Enteric-coated Mycophenolate Sodium (E-MPS, Myfortic) compared with Mycophenolate Mofetil (NNF) in de Novo Transplant Recipients: Results of a 12-Month, Single-blind, Randomized, Parallel-Group, Multicenter Study. The Journal of Heart and Lung Transplantation, Vol. 25, Number 8, 2006. **54.** Zackliczynski M et al. Letter: Elective conversion from CellCept to Myfortic under Control of Mycophenolate Acid Concentration in Stable Heart Transplant Recipients. He Journal of Heart and Lung Transplantation, Vol. 26, No. 3, 2007. **55.** Pharmacokinetics and variability of mycophenolic acid form enteric-coated mycophenolate sodium compared with mycophenolate mofetil in de novo heart transplant recipients. Clin Transplant 2997; 21: 18-23. **56.** Dumortier J et al. Conversion from Mycophenolate Mofetil to Enteric-coated Mycophenolate Sodium in Liver Transplant Patients Gastrointestinal Disorders: A Pilot Study. Liver Transplantation 12: 1342-1346, 2006. **57.** Cantisani GPC. Enteric-coated Mycophenolate Sodium Experience in Liver Transplant Patients. Transplantation Proceedings, 38: 932-933, 2006. **58.** Sollinger HW. Mycophenolate in transplantation. Clin Transplant 2004;18: 485-492. **59.** Chan L et al. Patient-Reported Gastrointestinal Symptom Burden and Health-Related Quality of Life following Conversion from Mycophenolate Mofetil to Enteric-Coated Mycophenolate Sodium. Transplantation Vol 81, No 9 2006. **60. Trapianto di rene.** Puig JM et al. Comparison of Mycophenolate Mofetil (MMF) and enteric-coated mycophenolate sodium (EC-MPS) pharmacokinetic profile in stable renal transplant (RT) patients treated with tacrolimus (FK) without steroids. American Journal of Transplantation 7: 349-349; 780 Suppl. 2 May 2007. Bolin P et al. Improvement in 3-Month Patient Reported Gastrointestinal Symptoms After Conversion From Mycophenolate Mofetil to Enteric-Coated Mycophenolate Sodium in Renal Transplant Patients. Transplantation 2007; 84: 1443-1451. Budde K et al. Conversion from Mycophenolate Mofetil to Enteric-Coated Mycophenolate Sodium in Maintenance Renal. Transplant Recipients Receiving Tacrolimus: Clinical Pharmacokinetics, and Pharmacodynamic Outcomes Transplantations 2007;83:417-424. **61. Trapianto di fegato.** Wiesner R et al. The Rapamune Liver Transplant Study Group. Sirolimus immunotherapy results in reduced rates of acute rejection in de novo orthotopic liver transplant recipients. Am J Transplantation 2002; Washington; 2002. Maheshwari A et al. Sirolimus Monotherapy Versus Sirolimus in Combination with Steroids and/or MMF for Immunosuppression After Liver transplantation. Dig Dis Sci 2006. Zagha H et

al. A comparison of sirolimus vs. calcineurin inhibitor-based immunosuppressive therapies in liver transplantation. *Alimentary Pharmacology and Therapeutics* 2006; 23 (4): 513-520. Trotter JF et al. Low incidence of cytomegalovirus disease in liver transplant recipients receiving sirolimus primary immunosuppression with 3-day corticosteroid taper. *Transplant Infect Dis* 2003; 5 (4): 174-180. Trotter JF. Sirolimus in liver transplantation. *Transplant Proc* 2003; 35 (3 Suppl): S193-200. Neff GW et al. Ten years of sirolimus therapy in orthotopic liver transplant recipients. *Transplant Proc* 2003; 35 (3 Suppl): S209-16. McAlister VC et al. Liver and kidney pancreas transplantation using tacrolimus, sirolimus and steroid immunosuppression. *Transplantation* 1999; 67 (9): S601. Watson CJ et al. Sirolimus: a potent new immunosuppressant for liver transplantation. *Transplantation* 1999;67(4):505-509. Kneteman NM et al. Sirolimus-based immunosuppression for liver transplantation in the presence of extended criteria for hepatocellular carcinoma. *Liver Transpl* 2004;10(10):1301. Neff GW et al. Sirolimus therapy in orthotopic liver transplant recipients with calcineurin inhibitor related chronic renal insufficiency. *Transplant Proc* 2003;35(8):3029-3031. Nair Set al. Sirolimus monotherapy in nephrotoxicity due to calcineurin inhibitors in liver transplant recipients. *Liver Transpl* 2003; 9 (2): 126-9. Nour B et al. Safety and effectiveness of conversion to sirolimus in liver transplant recipients with renal dysfunction. *Am J of Transplantation* 2002. Sanchez EQ et al. Sirolimus conversion after liver transplantation: improvement in measured glomerular filtration rate after 2 years. *Transplant Proc* 2005; 37 (10): 4416-23. Sindhi R et al. Replacing calcineurin inhibitors with mTOR inhibitors in children. *Pediatric Transplantation* 2005; 9 (3): 391-7. Vester U et al. Sirolimus rescue of renal failure in children after combined liver-kidney transplantation. *Pediatric Nephrology* 2005; 20 (5): 686-9. Casas-Melley AT et al. Improvement in renal function and rejection control in pediatric liver transplant recipients with the introduction of sirolimus. *Pediatr Transplant* 2004; 8 (4): 362-6. **62. Trapianti pediatrici di rene e fegato.** Sindhi R, Seward J, Mazariegos G, Soltys K, Seward L, Smith A, et al. Replacing calcineurin inhibitors with mTOR inhibitors in children. *Pediatric Transplantation* 2005;9(3):391-397. Vester U, Kranz B, Nadalin S, Paul A, Becker J, Hoyer PF. Sirolimus rescue of renal failure in children after combined liver-kidney transplantation. *Pediatric Nephrology* 2005;20(5):686-9. Casas-Melley AT, Falkenstein KP, Flynn LM, Ziegler VL, Dunn SP. Improvement in renal function and rejection control in pediatric liver transplant recipients with the introduction of sirolimus. *Pediatr Transplant* 2004;8(4):362-6. Markiewicz M, Kalicinski P, Teisseyre J, Ismail H, Kaminski A, Teisseyre M. Rapamycin in children after liver transplantation. *Transplant Proc* 2003;35(6):2284-6. **63. Trapianti di midollo.** (GVHD-Stem cell transplantation-malignancy-pediatric). Cutler C, Antin JH. Sirolimus for GVHD prophylaxis in allogeneic stem cell transplantation. *Bone Marrow Transplant* 2004;34(6):471-6. Rapamycin (sirolimus) for treatment of chro-

ic graft-versus-host disease. *Biol Blood Marrow Transplant.* 2005; 11: 1: 47-55. Marty FM, Lowry CM, Cutler CS, Campbell BJ, Fiumara K, Baden LR, et al. Voriconazole and Sirolimus Coadministration after Allogeneic Hematopoietic Stem Cell Transplantation. *Biol Blood Marrow Transplant* 2006;12(5):552-559. **64. Trapianto di pancreas, cuore, polmone.** Toso C, Baertschiger R, Morel P, Bosco D, Armanet M, Wojtuszczyzn A, et al. Sequential kidney/islet transplantation: efficacy and safety assessment of a steroid-free immunosuppression protocol. *Am J Transplant* 2006;6(5):1049-58. Knight RJ, Kerman RH, Zela S, Podbielski J, Podder H, Van Buren CT, et al. Pancreas transplantation utilizing thymoglobulin, sirolimus, and cyclosporine. *Transplantation* 2006;81(8):1101-5. Gleissner CA, Doesch A, Ehlermann P, Koch A, Sack FU, Katus HA, et al. Cyclosporine Withdrawal Improves Renal Function in Heart Transplant Patients on Reduced-Dose Cyclosporine Therapy. *Am J Transplant* 2006. Vazquez de Prada JA, Vilchez FG, Cobo M, Ruisanchez C, Valls MF, Ruano J, et al. Sirolimus in de novo heart transplant recipients with severe renal impairment. *Transplant International* 2006;19(3):245-248. Bestetti R, Theodoropoulos TA, Burdmann EA, Filho MA, Cordeiro JA, Villafanha D. Switch from Calcineurin Inhibitors to Sirolimus-Induced Renal Recovery in Heart Transplant Recipients in the Midterm Follow-up. *Transplantation* 2006;81(5):692-696. Laporta Hernandez R, Ussetti Gil P, Garcia Gallo C, de Pablo Gafas A, Carreno Hernandez MC, Ferreiro Alvarez MJ. Rapamycin in Lung Transplantation. *Transplant Proc* 2005;37(9):3999-4000. Sanchez EQ, Martin AP, Ikegami T, Uemura T, Narasimhan G, Goldstein RM, et al. Sirolimus conversion after liver transplantation: improvement in measured glomerular filtration rate after 2 years. *Transplant Proc* 2005;37(10):4416-23. **65. Trapianto di isole di Langerhans.** Shapiro AMJ, Lakey JRT, Ryan EA, Korbutt GS, Toth EL, Warnock GL, Kneteman NM, Rajotte RV: Islet transplantation in seven patients with type 1 diabetes mellitus using a glucocorticoid-free immunosuppressive regimen. *N Engl J Med* ; 343:230-238. Shapiro AMJ, Ricordi C, Hering BJ et al: International trial of the Edmonton Protocol for islet transplantation. *N Engl J Med* 2006; 355: 1318-30. Froud T, Baidal DA, Ponte G, Ferreira DV, Ricordi C, Alejandro R: Resolution of neurotoxicity and beta-cell toxicity in an islet transplant recipient following substitution of tacrolimus with MMF. *Cell transplant* 2006, 15 (7): 613-629. **66. Profilassi Tx polmone:** Treede et Al, 3rd ICI San Diego, US, 2004; abstract 22; Keenan et Al, *Ann Thoracic Surg* 1995,60:580; Treede et Al, *J Heart Lung Transplant* 2001; 20:511. **67. Profilassi Tx Rene pancreas e pancreas:** Bechstein et Al, *Transplantation* 2004;7:1221; J Malaise et Al and EUROSPK Study Group, *Transplantation Proceedings* 2005 37,2843-2845; F Saudek and the SPK Study Group, *Nephrology Dialysis Transplantation* 2005. **68. Profilassi Tx Intestino-Multiviscerale:** Abu Elmagd et Al, *Ann Surg* 2001;234:404. **69. Trapianto di Midollo:** Koehler MT et Al *Bone Marrow Transplantation* 1995 15:895; Kananaru A et Al for the

Japanese FK 506 Study Group; Bone Marrow Transplantation 1995 15:885; Nash R.A et al Blood 1995 85: 3746; Nash R.A et al Blood 1996 88:3634; Nash R.A et al Blood 2000 96:2062-2068.

70. Trapianto di isole di Langerhans: Shapiro AMJ, Lakey JRT, Ryan EA, Korbutt GS, Toth EL, Warnock GL, Kneteman NM, Rajotte RV: Islet transplantation in seven patients with type 1 diabetes mellitus using a glucocorticoid-free immunosuppressive regimen. *N Engl J Med* ; 343:230-238. Shapiro AMJ, Ricordi C, Hering BJ et al: International trial of the Edmonton Protocol for islet transplantation. *N Engl J Med* 2006; 355: 1318-30.

71. Generale: Undre N., *Nephrol Dial Transplant* (2003) 18 [Suppl 1]: i12-i15; Andreoni KA et al., *American Journal of Transplantation* 2007; 7 (Part 2): 1359-1375; Guerra G. et al., *Transplant International* ISSN 0934-0874; Webster A. et al., *BMJ*, doi:10.1136/bmj.38569.471007.

Micofenolato Mofetile (MMF): Vitko S. et al., *Transplantation* 2005;80: 1734-1741; Squifflet JP et al., *Transplantation Proceedings*, 34, 1584-1586 (2002); Rostaing L. et al., *Transplantation - Volume 79*, Number 7, April 15, 2005; Mc Taggart S., *NEPHROLOGY* 2007; 12, S106-S110; Augustine j.j. et al., *Transplantation* 2006;81: 1004-1009; H.T. Silva et al., *American Journal of Transplantation* 2007; 7: 595-608; Ekberg H. et al., *N Engl J Med* 2007;357:2562-75; McAlister H. et al., *Cochrane Database of Systematic Reviews* 2006, Issue 4. Art. No.: CD005161. DOI: 10.1002/14651858.CD005161.pub2; Pageaux GP. Et al., *LIVER TRANSPLANTATION* 12:1755-1760, 2006; Grimm M. et al., *American Journal of Transplantation* 2006; 6: 1387-1397; Kobashigawa JA et al., *American Journal of Transplantation* 2006; 6: 1377-1386; Bravo C. et al., *Transplantation Proceedings*, 39, 2409-2412 (2007); Gallon LG et al., *Transplantation* 2007;83: 1324-1329.

Acido Micofenolico: Budde K. Et al., *Transplantation* 2007;83: 417-424.

Sirolimus: Vitko S. et al., *Transplantation* 2005;80: 1734-1741; ; Augustine j.j. et al., *Transplantation* 2006;81: 1004-1009; Kobashigawa JA et al., *American Journal of Transplantation* 2006; 6: 1377-1386; Ciancio G. et al., *Transplantation* 2006;81: 845-852.

Everolimus: Chan L. et al., *Transplantation* 2008;85: 821-826; Kovarik JM et al., *Transplantation Proceedings*, 38, 3456-3458 (2006).

Azatioprina: Grimm M. et al., *American Journal of Transplantation* 2006; 6: 1387-1397; Bravo C. et al., *Transplantation Proceedings*, 39, 2409-2412 (2007); Jessupp M et al., *Curr. Opin.in organ Transplantation* 12: 536-542 (2007).

Anticorpi (mono o policlonali): Boillot O. et al., *Liver Transplantation*, Vol 11, No 1 (January), 2005: pp 61-67; H.T. Silva et al., *American Journal of Transplantation* 2007; 7: 595-608; Kobashigawa JA et al., *American Journal of Transplantation* 2006; 6: 1377-1386; Rostaing L. et al., *Transplantation - Volume 79*, Number 7, April 15, 2005; Bravo C. et al., *Transplantation Proceedings*, 39, 2409-2412 (2007); Maudgil A et al., *Pediatr. Drugs* 2007 9(5).

72. Once-Daily Intravenous Busulfan Given with Fludarabine as Conditioning for Allogeneic Stem Cell Transplantation: Study of Pharmacokinetics and Early Clinical Outcomes J. A. Russell, H. T. Tran, D. Quinlan, A.

Chaudhry, P. Duggan, C. Brown, D. Stewart, J. D. Ruether, D. Morris, S. Glöck, E. Gyonyor, B. S. Andersson *Biology of Blood and Marrow Transplantation* 8:468-476 (2002).

73. Graft-versus-host disease following allogeneic transplantation from HLA-identical sibling with antithymocyte globulin-based reduced-intensity preparative regimen Mohamad Mohty, Jacques-Olivier Bay, Catherine Faucher, Bachra Choufi, Karim Bilger, Olivier Tournilhac, Norbert Vey, Anne-Marie Stoppa, Diane Coso, Christian Chabannon, Patrice Viens, Dominique Maraninchi, and Didier Blaise. *Blood*, 15 July 2003 - Vol 102, Number 2.

74. Antithymocyte globulin affects the occurrence of acute and chronic graft-versus-host disease after a reduced-intensity conditioning regimen by modulating mixed chimerism induction and immune reconstitution. Nakai K, Mineishi S, Kami M, Saito T, Hori A, Kojima R, Imataki O, Hamaki T, Yoshihara S, Ohnishi M, Kim SW, Ando T, Fumitoh A, Kanda Y, Makimoto A, Tanosaki R, Kanai S, Heike Y, Ohnishi T, Kawano Y, Wakasugi H, Takaue Y. *Transplantation* 2003 Jun 27;75(12):2135-43.

75. Reduced-intensity preparative regimen and allogeneic stem cell transplantation for advanced solid tumors Blaise D, Bay JO, Faucher C, Michallet M, Boiron JM, Choufi B, Cahn JY, Gratecos N, Sotto JJ, François S, Fleury J, Mohty M, Chabannon C, Bilger K, Gravis G, Viret F, Braud AC, Bardou VJ, Maraninchi D, Viens P. *Blood* 2004 Jan 15;103(2):435-41.

76. Protective Conditioning for Acute Graft-versus-Host Disease Robert Lowsky, M.D., Tsuyoshi Takahashi, M.D., Ph.D., Ying Ping Liu, M.D., Sussan Dejbakhsh-Jones, M.S., F. Carl Grumet, M.D., Judith A. Shizuru, M.D., Ph.D., Ginna G. Laport, M.D., Keith E. Stockerl-Goldstein, M.D., Laura J. Johnston, M.D., Richard T. Hoppe, M.D., Daniel A. Bloch, Ph.D., Karl G. Blume, M.D., Robert S. Negrin, M.D., and Samuel Strober, M.D. *N Engl J Med* Sept 29, 2005 353:13.

77. Reduced incidence of Acute and Chronic Graft-versus-Host Disease with the Addition of Thymoglobulin to a Targeted Busulfan/Cyclophosphamide Regimen H. Joachim Deeg, Barry E. Storer, Michael Boeckh, Paul J. Martin, Jeannine S. McCune, David Myerson, Shelly Heimfeld, Mary E. Flowers, Claudio Anasetti, Kristine C. Doney, John A. Hansen, Hans-Peter Kiem, Richard A. Nash, Paul V. O'Donnell, Jerald P. Radich, Brenda M. Sandmaier, Bart L. Scott, Mohamed L. Sorror, E. Houston Warren, Robert P. Witherspoon, Ann Woolfrey, Frederick R. Appelbaum,ainer Storb. *Biology of Blood and Marrow Transplantation* 12:573-584 (2006).

78. Allogeneic transplantation for adult acute leukemia in first and second remission with a novel regimen incorporating daily intravenous busulfan, fludarabine, 400 CGY total-body irradiation, and Thymoglobulin Russell JA, Savoie ML, Balogh A, Turner AR, Larratt L, Chaudhry MA, Storek J, Bahlis NJ, Brown CB, Quinlan D, Geddes M, Stewart DA. *Biol Blood Marrow Transplant.* 2007 Jul;13(7):814-21.

79. Adult Recipients of Matched Related Donor Blood Cell Transplants Given Myeloablative Regimens Including Pretransplant Antithymocyte Globulin Have Lower Mortality Related to Graft-versus-Host Disease: A Matched

Pair Analysis James A. Russell, A. Robert Turner, Loree Larratt, Ahsan Chaudhry, Donald Morris, Christopher Brown,^{1,2} Diana Quinlan, Douglas Stewart Biology of Blood and Marrow Transplantation 13:299-306 (2007). **80. BMT** Winston DJ et al. Pharmacokinetics of ganciclovir after oral valganciclovir versus intravenous ganciclovir in allogeneic stem cell transplant patients with graft-versus-host disease of the gastrointestinal tract. Biol Blood Marrow Transplant 2006; 12 (6): 635-40. *The pharmacokinetics of ganciclovir after oral valganciclovir versus intravenous ganciclovir were compared in valganciclovir could be a useful alternative to intravenous ganciclovir in certain stable stem cell transplant patients who require prophylaxis or pre-emptive therapy for cytomegalovirus infection.* Ayala E et al. Valganciclovir is safe and effective as pre-emptive therapy for CMV infection in allogeneic hematopoietic stem cell transplantation. Bone Marrow Transplant 2006; 37 (9): 851-6 *Pre-emptive therapy of CMV infection with oral VGC is safe and effective in allogeneic HSCT recipients.* Van der Heiden PLJ et al. Oral valganciclovir as pre-emptive therapy has similar efficacy on cytomegalovirus DNA load reduction as intravenous ganciclovir in allogeneic stem cell transplantation recipients. Bone Marrow Transplant 2006; 37 (7): 693-8. *The efficacy and safety of oral valganciclovir was compared to ganciclovir i.v. in pre-emptive treatment of cytomegalovirus (CMV) in T-cell-depleted allogeneic stem cell transplant (allo-SCT) recipients. In conclusion, pre-emptive treatment with valganciclovir and ganciclovir, led to similar reduction of CMV DNA load. Oral valganciclovir is an attractive and safe alternative for pre-emptive CMV treatment in T-cell-depleted allo-SCT recipients.* Einsele H et al. Oral valganciclovir leads to

higher exposure to ganciclovir than intravenous ganciclovir in patients following allogeneic stem cell transplantation. Blood 2006; 107 (7): 3002-8. *This supports the use of V-GCV in SCT, even in patients with I-GVHD grades I-II. Due to higher exposure after VGCV compared with IV-GCV, patients should be monitored carefully for safety reasons.*

Boeckh M et al. Cytomegalovirus in hematopoietic stem cell transplant recipients: Current status, known challenges, and future strategies. Biol Blood Marrow Transplant 2003; 9 (9): 543-58. *Strategies currently being investigated include long-term suppression of CMV with valganciclovir for the prevention of late CMV infection and disease, adoptive transfer of CMV-specific T cells, and donor and recipient vaccination strategies.*

81. Infezione da Cytomegalovirus in pazienti pediatrici con trapianto. Clark BS et al. Valganciclovir for the prophylaxis of cytomegalovirus disease in pediatric liver transplant recipients. Transplantation. 2004; 77 (9): 1480

Burri M et al. Oral valganciclovir in children: single dose pharmacokinetics in a six-year-old girl. Ped Infectious Dis J 2004; 23 (3): 263-266. Reasonable RR, Paya CV. Valganciclovir for the prevention and treatment of cytomegalovirus disease in immunocompromised hosts. Expert Rev. Anti-infect. Ther. 2004; 2 (1): 27-42) Vaudry W et al. Safety of oral valganciclovir powder for oral solution in pediatric solid organ transplant recipients. Am J Transplant / Transplantation 2006 WTC 215 Abs 441. Bouw R et al. Ganciclovir pharmacokinetics in pediatric solid organ transplant recipients after administration of valganciclovir oral solution. Am J Transplant / Transplantation 2006 WTC 215 Abs 442.